

Mini-dark halos with intermediate mass black holes

HongSheng Zhao and Joseph Silk

(Dated: 1 June 2005 on Phys. Rev. Letters 95, 011301)

We argue that the Milky Way (MW) contains thousands intermediate mass black holes (IMBHs) and minihalos with a fraction of IMBHs still being enshrouded in extremely dense mini-spikes of dark matter (DM) particles. Each containing $10^6 M_\odot$ of dark particles and no baryons in a sphere of 50 pc radius, the minihalos are dense enough to survive the Milky Way tide with the nearest minihalo within 2 kpc from the Sun. The IMBH is formed off-centre in a minihalo by gas accretion, and its growth adiabatically compresses a finite density of surrounding dark matter into a $r^{-1.5}$ mini-spike. Some IMBHs recentre on their minihalos after dynamical friction, and some IMBHs are ejected by birth kicks. Detectable by GLAST, the mini-spikes and minihalos should stand out the background and dominate the neutralino annihilation in the smooth MW and satellite galaxies. If they are the unidentified EGRET sources, upper limits can be set on the branching ratio of neutralino annihilation. The supermassive BH of the MW, if originates from an IMBH, is also likely enshrouded with a mini-spike.

PACS numbers: 95.35.+d; 98.35.Gi; 98.35.Jk

Supermassive black holes (SMBH) likely grow from smaller intermediate mass black holes (IMBH) by gas accretion; to be massive enough and early enough, a seed of $\sim 1000 M_\odot$ at $z \sim 20$ is necessary to grow to a $3 \times 10^9 M_\odot$ SMBH at $z = 6$ via Eddington accretion of gas (e.g., [1, 2] and references therein). Gas accretion is favoured because of the empirical result [3, 4] that requires radiatively efficient accretion in order to account for both the quasar luminosity function and the mass in black holes at present. Conversely, once the quasar/AGN phase is over, black holes undergo radiatively inefficient accretion as is the case for Sgr A* (e.g., [5]). Here we explore implications of growth of SMBH by accretion in the protogalactic environment. We will study the evolution of the seed IMBHs. These smaller IMBHs are plausibly formed in a subset of the first objects, where accretion rates are high. Hierarchical merging of minihalos leads to a population of IMBHs in MW-sized halos. A mini-spike of CDM is inevitable around the IMBH due to adiabatic growth in the minihalo. The mini-spike is robust to halo merging, and survives around IMBHs that populate the halo. The subsequent role of BH merging depends on dynamical friction and loss cone refilling. These processes are likely to nurture the growth of the central seed IMBH into a supermassive BH.

The IMBHs, and in particular, the dark matter spikes, are potentially observable via neutralino annihilations. We mention two specific examples. Firstly, a central IMBH is inferred to be associated with a group of massive stars IRS 13 offset by about 2 arcsec from the Galactic centre (SgrA*) [6]. Secondly, the Whipple/HESS detection of > 3 TeV gamma rays from the Galactic centre ([7, 8, 9]) would require a high annihilation cross section ~ 10 pb if this emission is due to a neutralino annihilation signal produced by an NFW[10] halo profile [11]. Unfortunately, to create the Λ CDM relic density requires a cross-section of ≤ 1 pb. Nor even for subdominant neutralinos can SUSY theory easily produce such a large cross-section above 10 TeV. A spike may provide the best way of reconciling the high annihilation flux with the dark matter relic abundance [11, 12].

The redshift and mass of the first clouds, stars and IMBHs: The temperature T of the first clouds that contain gravitationally bound baryons depends mostly on the virial mass M with $M = 10^6 M_\odot \left[\frac{T}{10^3 \text{K}} \frac{10}{1+z} \right]^{3/2}$, where we have scaled with typical values of the redshift z and temperature T ; cooling by H_2 requires a virial temperature in excess of several hundred degrees K. The collapse redshift for clouds of mass M is $\approx 0.6\nu \times 16(M/10^6 M_\odot)^\epsilon$, where $\epsilon \ll 1$ and ν is the height of the gaussian peak. Hence the desired redshift is sensitive primarily to the peak rarity, and must be in the range 10-30. The WMAP-inferred optical depth of the universe to Thomson scattering also suggests a redshift of 15-20 for the first objects that reionised the universe.

The mass of the first objects can be estimated by noting that the accretion rate is of order $v_s^3/G \sim 10^{-4} - 10^{-3} M_\odot/\text{yr}$, where we take v_s to be the virial velocity. Accretion can only be halted by feedback, and it takes at least 10^6 years to generate feedback from massive stars, hence the masses of the first stars are of order $100 - 1000 M_\odot$, as found in numerical simulations of primordial gas clouds by [13]. Finally, in clouds of 10^4K where Lyman alpha cooling dominates, the accretion rate onto the protostellar core is around $0.03 M_\odot/\text{yr}$, hence feedback cannot inhibit the formation of an IMBH, of mass up to $\sim 10^4 M_\odot$.

Annihilation flux of (mini-)halo models with finite core and plateau: The (bolometric) annihilation flux integrated over a sphere of WIMPs of mass m_χ and number density $n(r) = \frac{\rho(r)}{m_\chi}$ at radius r is

$$L_{\chi\bar{\chi}}(< r) = \int_0^r 4\pi r^2 dr \langle \sigma v \rangle m_\chi c^2 \left[\frac{\rho(r)}{m_\chi}, \frac{\rho(r_p)}{m_\chi} \right]_{\min}^2, \quad (1)$$

where annihilation imposes a plateau of size $r < r_p$ on the present-day dark matter density because neutralino number density above $\frac{\rho(r_p)}{m_\chi} \equiv \frac{H_0}{\langle \sigma v \rangle} \sim \frac{10^8 M_\odot \text{pc}^{-3}}{50 \text{GeV}} \sim 10^8 \text{cm}^{-3}$ will annihilate away over a Hubble time (e.g., [14, 15]), where the cross-section $\langle \sigma v \rangle / c \sim 1$ pb, fixed by the relic abundance Ω_χ required in a Λ CDM cosmology. The total detectable flux

quickly converges at large radius since $\rho(r)$ generally drops as a steeper than $r^{-1.5}$ power-law beyond r_p . Adopting the typical SUSY mass $m_\chi = 50\text{GeV}$, we have

$$\frac{L_{\chi\bar{\chi}}(< r)}{2 \times 10^3 L_\odot} = \int_0^r \left[1, \frac{\rho(r)}{10^8 \text{M}_\odot \text{pc}^{-3}} \right]_{\min}^2 \frac{r^2 dr}{(10^{-3} \text{pc})^3}. \quad (2)$$

Note that less than 10% of the above calculated flux for neutralino models will likely go into photons (γ -rays). More generally and crudely speaking, $L_{\chi\bar{\chi}}(< r) \propto m_\chi^{-1}$.

We model the density $\rho(r)$ of the minihalos and the MW halo with a subset of models of Zhao[16] (also known as generalized-NFW profiles). To be *conservative in estimating the annihilation signal*, we choose a subset of initial halo models with a *finite core and without any outer truncation*. Our models are also inspired by the recent exponential-like model of Navarro et al. ([17], N04 model hereafter), but are *easier to work with analytically*. Specifically, we model halos with density at radius r given by

$$\rho(r) = \frac{\rho(r_c)}{[x/2 + 1/2]^{12}}, \quad x \equiv \left(\frac{r}{r_c} \right)^{\frac{1}{3}}. \quad (3)$$

Here r_c is, as in an NFW model, where the local density power-law index $d \log \rho / d \log r = -2$. Our models are specified by two parameters r_c and $\rho_c \equiv \rho(r_c)$, and the virial radius $r_{\text{vir}} \equiv cr_c$ is given implicitly by

$$\frac{3M_{\text{vir}}}{4\pi r_{\text{vir}}^3} = 176\rho_{\text{crit}}(z) = \frac{2\rho(r_c)(1 + c^{\frac{1}{3}}/5 + c^{\frac{2}{3}}/55)}{(1/2 + c^{\frac{1}{3}}/2)^{11}}, \quad (4)$$

i.e., the mean density $\bar{\rho}(r_{\text{vir}}) = \frac{3M_{\text{vir}}}{4\pi r_{\text{vir}}^3}$ is approximately 176 times the critical density $\rho_{\text{crit}}(z)$ of the universe at a given formation redshift z . Note the density ratio $\bar{\rho}(r_c)/\rho(r_c) = 2.436, 2.3178$ and 2.4644 for our cored model, NFW model[10], and N04 model. Unlike the NFW model, our model has an outer r^{-4} density profile without truncation, and a finite central density and a finite asymptotic mass. For models with the same core radius and density normalisation $\rho(r_c)$, $M(r_{\text{vir}})$ differs by only 10 percent between NFW models and our model for typical concentration $c = 1 - 1000$.

There are a few advantages of our model. It is straightforward to derive and show that the potential, the circular velocity curve and even the velocity dispersion curve of this class of halo models are all *simple analytical* rational functions of $r^{1/3}$, while some special functions enter the NFW and N04 models. As shown in Fig.1, our model reproduces the density, mass, velocities of NFW and N04 very well over the range $0.01r_c < r < 100r_c$ at least for systems with $c \geq 1$.

Formation of minihalos and massive black holes: Current numerical simulations suggest that the first stars form from H_2 clouds in dark minihalos of mass about $3 \times 10^5 h^{-1} \text{M}_\odot$, virial radius of $60 h^{-1} \text{pc}$ and virial velocity of 5 km s^{-1} , and initial virial temperature of 1000 Kelvin (for a molecular weight of unity) at a redshift of $z \sim 20$, when the universe was about 0.15 Gyr old. The cooling time is comparable to the dynamical time, both being of order 1 Myr, and




FIG. 1: log-log radial profiles of the rescaled halo density $\rho(r)/\rho(r_c)$, escape velocity $V_{\text{esc}}(r)/(K\sqrt{2})$, dispersion $2\sigma(r)/K$, circular velocity $V_{\text{circ}}(r)/K$, and enclosed mass $M(r)/100\rho(r_c)r_c^3$ in our cored halo model (crosses), NFW model (blue dashed lines) and N04 model (red lines). Here $K = \sqrt{G\rho(r_c)r_c^2}$, and r_c is the core radius.

self-gravitating isothermal clouds of order a thousand solar masses are able to develop. Once the H_2 metal-free gas in these systems cools to a temperature of $\sim 200\text{K}$ and density of $\sim 10^4 \text{cm}^{-3}$, the H_2 cooling saturates and subfragmentation ceases [18]. These clouds are generally not at the exact centres of the minihalos, but have dynamically decayed to the central parsec of the minihalos with typical random motions of less than 10 km s^{-1} . The first stars are generally massive (about $100 - 1000 \text{M}_\odot$) because of inefficient cooling with zero metals. They are short-lived. According to [19] (see their Fig.2), these very massive zero-metallicity stars have large helium cores, which are unstable to pulsations due to electron-position pair instability. A supernovae leaving no remnant happens for $140 - 260 \text{M}_\odot$ stars, and a more massive star collapses into an IMBH after losing about half of the initial mass by wind. However, Lyman alpha-cooling clouds would be expected to form more massive IMBHs, up to $10^3 - 10^4 \text{M}_\odot$.

There are several plausible outcomes for the IMBH and the gas inside a minihalo. In some cases, the IMBH could be ejected out of the shallow potential well of the minihalo if born with a significant kick/recoil velocity (greater than the 30 km s^{-1} escape velocity at the centre of the minihalo) [20]. In other cases, the IMBH spirals to the centre of the mini-halo due to dynamical friction (the timescale $t_{\text{df}} \sim (r/\text{pc})^2 \sim 1 \text{ Myr}$ for a thousand solar mass BH formed at 1 pc). At the centre or en route, these IMBHs may accrete gas at a super-Eddington rate with a mass-doubling time scale $400\epsilon \ln(2) \approx 1 - 20 \text{ Myr}$ for a plausible range of the accretion efficiency $\epsilon = 0.005 - 0.1$. Eventually after some uncertain

period of order 10 Myr, the first SNe progenitors are born in these minihalos, and shortly afterwards (about 4 Myr) most of the gas in the minihalos (perhaps with the exception of the tightly bound accretion disk around the IMBH) is cleared out when the SNe explode. Thus the only remnants of the first generation of star formation are a population of minihalos and IMBHs, which are either well-centred due to dynamical friction or well-separated due to strong kicks. To be specific, we consider minihalos specified by eq. (3) with $r_c = 1.6\text{pc}$ and $\rho(r_c) = 500\text{M}_\odot\text{pc}^{-3}$ (comparable to an NFW halo of $r_{\text{vir}} = 50 \times r_c = 80\text{pc}$). We assume conservatively that 10 percent of the minihalos harbour IMBHs which eventually grow to a final mass of $10^2\text{M}_\odot$ or $10^3\text{M}_\odot$. These values are plausible for a r.m.s. kick velocity $\sim 100\text{km s}^{-1}$, and accretion efficiency $\epsilon \sim 0.02$ in the mini-halo in 10 Myrs. So there are of order 10^3 minihalos containing IMBHs.

Adiabatic compression of dark matter in minihalos: Now consider the effect on the dark matter. The formation and the growth of the IMBH will compress the surrounding dark matter. The compression is close to adiabatic since the formation of the IMBH and the subsequent accretion are on time scales of the cooling time and Salpeter time; both are somewhat longer than the dynamical time. Given the finite density and phase space density of dark matter at an off-centre position, we expect a moderate $r^{-3/2}$ power-law dark matter density distribution surrounding the IMBH. This type of cusp (or "spike") is the result of growth in a uniform background[25]; a steeper cusp occurs if a well-centred BH grows adiabatically (e.g., [15]). Such a mini-spike will survive dynamical friction, and be added to the central core of the minihalo once the IMBH re-centers. We expect of order of $m_\bullet \sim 1000\text{M}_\odot$ of dark matter in this spike. So the dark matter mass profile with a mini-spike is

$$M^{\text{sp}}(r) = \left(1 + \frac{b^{3/2}}{r^{3/2}}\right) M(r), \quad M(b) = m_\bullet, \quad (5)$$

where $b = 0.045\text{pc}$ or 0.14pc so that $M^{\text{sp}}(b) = 2M(b) = m_\bullet = 10^2\text{M}_\odot$ or $10^3\text{M}_\odot$.

We consider several possibilities for the dark matter. Fig.2 shows the resulting dark halo density and the annihilation flux. In the simplest case, the minihalos have the pristine cored density profile; the IMBH either never formed or has been kicked out of the system. For a minihalo which keeps its IMBH, the IMBH draws in a mini-spike of dark matter with a plateau inside about 0.005pc depending on its mass. The flux per log radius bin is of order $10^4 L_\odot$, and most of it comes from within 0.01pc of the minihalos.

Observability of minihalos inside the MW: By a redshift of $z = 1 - 6$, nearly all the minihalos and IMBHs are bound inside galaxy-sized objects. About $10^3 - 10^4$ should be bound inside a Milky-Way-sized galaxy [21], which we assume to have a density given by eq. (3) with $r_c = 11\text{kpc}$ and $\rho(r_c) = 0.0055\text{M}_\odot\text{pc}^{-3}$ (this is consistent with the local DM density and resembles an NFW model of $r_{\text{vir}} = 8r_c$). These minihalos should be distributed in a way similar to the overall




FIG. 2: **Left:** log-log plot of the radial profiles of spherically averaged density (black lines) in $\text{M}_\odot\text{pc}^{-3}$ inside a mini-halo in three cases: a strong mini-spike around a $1000\text{M}_\odot$ IMBH (thick line), or a moderate mini-spike around a $100\text{M}_\odot$ IMBH, or a pure cored minihalo where the IMBH either was never formed or was ejected (thin line). Lines (green parallel) of constant enclosed mass $1\text{M}_\odot, 10\text{M}_\odot, \dots, 10^{12}\text{M}_\odot$ are shown to guide the eye. For comparison are the average densities inside a dwarf galaxy (lower thick line, with dispersion of 10km s^{-1} and a core of 200pc) and inside the pristine MW halo (lower right thin line). **Right:** the logarithm of the rescaled annihilation (bolometric) luminosity $\frac{L_{\chi\bar{\chi}}(<r)}{L_\odot} \times \frac{m_\chi}{50\text{GeV}}$ (nearly horizontal red lines to the right) of these models. Also shown are the total luminosity of 10^3 minihalos with $100\text{M}_\odot$ IMBHs (thickest black at top middle), the luminosity of the MW as a whole (thick red) or scaled down by a factor $\frac{\Delta\Omega}{4\pi}$ for a cone of $\Omega = 10^{-3}$ (thin red). The minihalos stand out as clumps above the MW background.

dark matter profile of the galaxy, producing a clumpy annihilation brightness distribution; we assume the stars and the central SMBH in the MW form later.

A typical nearby minihalo at about 3 kpc from the Sun sustains a solid angle $\Delta\Omega \sim 10^{-3}$ (about the resolution of EGRET) comparable to the angular size of dwarf galaxy satellites at 100kpc , but minihalos are much more luminous and brighter in annihilation than dwarf galaxies because of minihalo's much denser core than dwarf galaxies; the latter have central dark matter densities of $(0.05 - 0.5)\text{M}_\odot\text{pc}^{-3}$ [26]. Except towards the Galactic center a minihalo stands out in density and annihilation flux from the MW background in a beam of $\Delta\Omega \leq 10^{-3}$ at all distances (cf. Fig. 2). A population of 10^3 minihalos with $100\text{M}_\odot$ IMBHs is about 100 times more luminous than the MW background as a whole. These minihalos are low-mass ($\sim 10^6\text{M}_\odot$) but dense (about $1\text{M}_\odot\text{pc}^{-3}$ on a scale of 30pc) enough so that 99 percent of the minihalos survive intact in the halo, and follow the overall distribution of dark matter. There are of order 10 minihalos within 1 kpc of the MW centre, where dynamical friction could bring one of these minihalos to the centre of the MW within a Hubble time (cf. Fig.3 of [15]), without being slowed down by tidal

stripping [22]; note that these minihalos are typically denser than the MW halo on a scale of 60pc. If the IMBH of the minihalo could serve as a seed for the supermassive BH in the MW.

The isolated IMBHs are virtually invisible. The X-ray luminosity due to Bondi-Hoyle accretion of an IMBH passing through the MW gaseous disk is estimated to be $(\epsilon/0.1)(n/30\text{cm}^{-3})(m_\bullet/300M_\odot)^2L_\odot$ [23]. So the signal is much fainter than the million-solar-mass BHs of Lacey & Ostriker[23]. Such a low luminosity is just detectable with present facilities if the IMBH is close-by (say 2 kpc), but the chance of a BH passing through a cold H_2 or HI cloud (with filling factors 0.3% and 2% respectively) at a given time is very low. The local density of IMBHs is about 0.3 kpc^{-3} assuming 10^4 IMBHs in the whole halo, and we expect only of order one IMBH hitting a cloud with a filling factor of $\sim 1\%$ in the entire disk of 8 kpc in radius and 200pc in thickness, an approximation for the cold H_2 and HI cloud distribution. Also being much fewer than the million-solar-mass BHs of [23], the minihalos do no damage to the MW disk.

Despite being a dynamically subdominant population, the minihalos and IMBHs speed up two-body relaxation in the MW. The relaxation time is given by [23] $\frac{1\text{Myr}}{t_{\text{rlx}}} = \frac{\rho(r)\sigma^{-3}}{10^6M_\odot\text{pc}^{-3}(100\text{km s}^{-1})^{-3}} \frac{\bar{M}}{10^3M_\odot}$, where $\bar{M} = \frac{f_M \ln \Delta}{20} \frac{M^2 + m_\bullet^2}{M + m_\bullet} \sim 1 - 1000M_\odot$ is the mass density-weighted mean mass of deflectors depending on tidal stripping; the lower value is for minihalos well inside 1-10 kpc of the Galactic centre, and $f_M \sim 10^{-3}$ is the fraction of matter mass density in the form of minihalos of mass M . The heavier the deflectors, the faster the relaxation. So compared to the case of purely stellar deflectors, the relaxation is about $\bar{M}/M_\odot \sim 1 - 1000$ times more efficient. This might help to refill the loss-cone of the central BH by scattering stars and dark matter particles [24]. To refill the loss-cone of $3 \times 10^6 M_\odot$ BH on the scale of the 1pc would require a relaxation time of order a Hubble time.

Our model predicts around $1500 L_\odot$ (bolometric) per solar mass in spike neutralinos. Observationally, the annihilation signal is best seen via gamma rays. EGRET observations set upper limits on unidentified sources, in particular on the source near the Galactic Centre, in gamma rays above 1 GeV of around $100L_\odot$ [27]. For our models to be consistent with observational data in the MW and a gamma ray branching ratio of order 10 percent, we require of order a solar mass of the neutralinos in the spike for typical SUSY neutralinos

with $m_\chi = 50\text{GeV}$. This is consistent with our models of adiabatic growth of $10^2 M_\odot$ IMBHs and models with ejected IMBH. The data would also be consistent with the $10^3 M_\odot$ IMBH model in case of heavier neutralinos ($m_\chi \sim 10\text{TeV}$ [28]). Evidently the GLAST satellite, with two orders of magnitude more sensitivity than EGRET, will be capable of setting stringent constraints on halo IMBH.

We thank the referee for many detailed comments which help to improve the presentation here. HSZ acknowledges travel and publication support from Chinese NSF grant 10428308 and UK PPARC Advanced Fellowship.

-
- [1] van der Marel, R., 2003, Coevolution of black holes and galaxies, ed. L.C. Ho, Cambridge Univ. Press, astro-ph/0302101
 - [2] Gnedin O., 2001, Class. and Quan. Gravity, 18, 3983 (astro-ph/0108070)
 - [3] Soltan, A. 1982, MNRAS, 200, 115
 - [4] Yu, Q. and Tremaine, S. 2002, MNRAS, 335, 965
 - [5] Goldston, J., Quataert, E. and Igumenshchev, I. 2005, astr-ph/0411627, ApJ (in press)
 - [6] Maillard J.P., et al. 2004, A&A, 423, 155
 - [7] Kosack et al., 2004, ApJ, 608, L97,
 - [8] Aharonian et al. 2004, A&A, 425, L13
 - [9] Buckley J., 2005, Galactic Center News, vol. 20, March issue,
 - [10] Navarro J. F., Frenk C. S., White S. D. M., 1996, ApJ, 462, 563
 - [11] Silk, J. 2004, (astro-ph/0412297)
 - [12] Hooper, D. and March-Russell, J., 2004, hep-ph/0412048.
 - [13] Abel, T., Bryan, G. & Norman, M. 2002, Science, 295, 93
 - [14] Gondolo P. & Silk J. 1999, PhRvL., 83, 1719
 - [15] Ullio P., Zhao H., Kamionkowski M. 2001, PRD, 64, 3504
 - [16] Zhao H.S., 1996, MNRAS, 278, 488
 - [17] Navarro J. F., et al. 2004, ApJ, 349, 1039 (N04)
 - [18] Bromm, V. and Larson, R. 2004, ARAA, 42, 79
 - [19] Heger A. & Woosley S.E., 2002, ApJ, 567, 532
 - [20] Madau, P. and Quataert, E. 2004, ApJ, 606, L17
 - [21] Madau, P. and Rees, M. 2001, ApJ, 551, L27
 - [22] Zhao H.S., 2004, MNRAS, 351, 891
 - [23] Lacey C. & Ostriker J. , 1985, 299, 633
 - [24] Zhao H.S., Haehnelt M, Rees M., 2002, NewA., 7, 385
 - [25] Binney J.J. & Tremaine S. 1987, Galactic Dynamics, Princeton University Press.
 - [26] Mateo M. 1998, 36, 435, Annual Review of Astronomy & Astrophysics
 - [27] Nolan P. et al. 2003, , ApJ, 597, 615
 - [28] Bertone G. & Merritt D. 2005, astro-ph/0501555, PRL submitted